AD-A275 813

FINAL REPORT

GRANT#: N00014-89-J-1865

R&T CODE: 4414810

PRINCIPALINVESTIGATOR: Joseph F. Siebenaller

INSTITUTION: Louisiana State University

GRANTTITLE: Effects of Pressure on Membrane-AssociatedReceptors and Effector Elements

REPORTING PERIOD: 1 June 1993 - 30 November 1993

AWARD PERIOD: July 1989 - November 1993

<u>OBJECTIVE</u>: To identify the effects of moderate hydrostatic pressure on receptor and effector elements involved in transmembrane signal transduction, we examined the A_i adenosine receptor - inhibitory G protein (G_i) - adenylyl cyclase signal transduction complex. Our experiments were designed to identify and define at the molecular level pressure effects on system components in isolation and on the entire functional complex.

APPROACH: Two marine teleost species, Sevastolobus alascanus and S. altivelis which live at different depths were used as a model to study pressure adaptation. This approach of parallel comparisons has permitted the identification of patterns of pressure adaptation. Among the techniques employed were assays of adenylyl cyclase activity, assays of the high-affinity GTP as activity of G proteins, antibody quantitation of G proteins, pertussis and cholera toxin ADP-ribosylation of G protein subunits, affinity labeling and peptide mapping.

ACCOMPLISHMENTS: We have characterized the effects of hydrostatic pressure on the A. adenosine receptor - inhibitory G protein (G) - adenylyl cyclase signaling complex. Hydrostatic pressure inhibits basal adenylyl cyclase activity, increases the K_m of ATP for adenylyl cyclase and decreases the efficacy of agonists in inhibiting cAMP accumulation. Pressure also affects the high-affinity GTP ase activity in brain membrane preparations. There are two GTPase activities, one with a low- and one with a high-K of GTP. The high-affinity GTP ase activity, characteristic of the \alpha subunits of the guanine nucleotide binding regulatory protein (G protein) pool, was stimulated by the A, adenosine receptor and muscarinic cholinergic receptor agonists. High-affinity hydrolysis of GTP, measured at 0.3 µM GTP, was stimulated 22% in both species by 340 atm pressure. At 340 atm pressure, the apparent K_m of GTP is decreased approximately 10% in each of the species, and the V_{max} values are increased 11 and 15.9% in S. alascanus and S. altivelis, respectively. The apparent volume changes associated with the decreased K_ of GTP values and the increased V values ranged from -7.0 to -9.9 ml/mol. Increased pressure, however, markedly decreases the efficacy of adenosine and muscarinic agonists in stimulating GTPase activity. These results suggest that the effects of increased hydrostatic pressure on transmembrane signal transduction by this system may stem, at least in part, from pressure-increased GTP hydrolysis and the concomitant termination of inhibitory signal transduction.

Pertussis toxin-catalyzed [³²P]ADP-ribosylation at atmospheric pressure and 5°C was used to probe the guanine nucleotide binding regulatory proteins G₁ and G₂ in brain membranes from the two Sebastolobus species. The membranes from the deeper-living S. altivelis consistently incorporated more [³²P]ADP than the membranes from S. alascanus. Because the heterotrimeric holoprotein is the preferred substrate for the ribosylation reaction, the modulatory effects of the guanyl nucleotides GDP and GTPγS on ribosylation were assessed. GDP increased [³²P]ADP-ribosylation of the α subunits in S. altivelis. Only the highest concentration tested (1000 μM) increased [³²P]ADP-ribosylationin S. alascanus brain membranes to a slight extent. Increasing concentrations of GTPγS suppressed [³²P]ADP-ribosylation in S. alascanus brain membranes, presumably by promoting dissociation of the holotrimer. GTPγS had much less of an effect on the S. altivelis brain membranes. These differences in the extent of ADP-ribosylation and the modulatory effects of guanyl nucleotides may relate to the differential extent of coupling receptors to G

94

2

17

045

This accument has been approved for public release and sale; its distribution is unlimited.

DTIC QUALITY INSPECTED 2

94-05

proteins in these two species. The effects of hydrostatic pressure on the PTX-catalyzed[32P]ADP-ribosylation reaction differ in brain membranes of the two species. In the presence of GDP, the ribosylation reaction in S. altivelis is unaffected by pressures up to 360 atm. 68 atm of pressure inhibits ribosylation 40% in S. alascanus membranes.

SIGNIFICANCE: Increased hydrostatic pressure decreases basal adenylyl cyclase activity and decreases the efficacy of A adenosine receptor agonists in modulating adenylyl cyclase in these species. The effects of increased hydrostatic pressure on transmembrane signal transduction by this system stem, at least in part from pressure-increased GTP hydrolysis and the concomitant termination of inhibitory signal transduction, and the degree of coupling of the G protein pool to receptors. A general effect of pressure on agonist-receptor interactions with the G protein pool would be an important selective force in adaptation to the deep sea. Disruption of receptor coupling to effector elements via G proteins by pressure would be a critical impediment to colonization of the deep ocean by shallow-living organisms. Such a pressure-barrier to colonization would have important consequences both for the vertical distribution patterns of marine species, and the functioning of divers in this environment.

PUBLICATIONS AND ABSTRACTS:

- 1. J.F. Siebenaller and G.N. Somero (1989) Biochemical adaptation to the deep sea. Reviews in Aquatic Sciences 1: 1-27.
- 2. T.F. Murray, T.A. Blair, M. Leid, P.H. Franklin and J.F. Siebenaller (1989) A adenosine receptors in heart: Functional and biochemical consequences of activation. In: Probing Bioactive Mechanisms. P.S. Magee, D.R. Henry and J.H. Block (ed.), American Chemical Society Books, pp. 232-242.
- 3. J.F. Siebenaller and T. F. Murray (1990) A adenosine receptor modulation of adenylyl cyclase of a deep-living teleost fish, *Antimora rostrata*. Biological Bulletin 178: 65-73.
- 4. J.F. Siebenaller (1991) Pressure as an environmental variable: Magnitude and mechanisms of perturbation. In: *Biochemistry and Molecular Biology of Fishes*, Vol. 1, P.W. Hochachka and T.P. Mommsen (editors), Elsevier Press, pp. 323-343.
- 5. J.F. Siebenaller, A.F. Hagar and T.F. Murray (1991) The effects of hydrostatic pressure on A₁ adenosine receptor signal transduction in brain membranes of two congeneric marine fishes. Journal of Experimental Biology 159: 23-43.
- 6. B.J. Davis and J.F. Siebenaller (1992) Proteolysis at pressure and HPLC peptide mapping of M₄-LDH homologs from marine fishes living at different depths. International Journal of Biochemistry 24: 1135-1139.
- 7. J.F. Siebenaller and T.F. Murray (1994) The effects of hydrostatic pressure on the low-K GTPase in brain membranes from two congeneric marine fishes. Journal of Comparative Physiology B 164, in press.
- 8. J.F. Siebenaller and T.F. Murray (1993) Membrane preparation and receptor assays adenosine. In: *Biochemistry and Molecular Biology of Fishes*, Volume 3. P.W. Hochachka and T.P. Mommsen, editors, Elsevier Press, in press.
- 9. T.F. Murray and J.F. Siebena'ler (1994) Differential susceptibility of guanine nucleotide-binding proteins to pertussis toxin-catalyzed ADP-ribosylation in brain membranes of two congeneric marine fishes. Biological Bulletin 185: 346-354.
- 10. J.F. Siebenaller and T.F. Murray (1994) The effects of hydrostatic pressure on pertussis toxin-catalyzed ribosylation of guanine nucleotide-binding proteins from two congeneric marine fishes. CBP Part B, in press.
- 11. J.F. Siebenaller and T.F. Murray (1993) The effects of pressure on G protein-coupled signal transduction. In: *Biochemistry and Molecular Biology of Fishes*, Volume 5. P.W. Hochachka and T.P. Mommsen, editors, Elsevier Press, in press.
- 12. M. Leid, T.F. Murray, P.H. Franklin and J.F. Siebenaller (1989) A adenosine receptor regulation of adenylyl cyclase activity in brain membranes of a deep-living teleost fish. Society for Neuroscience Abstracts, Volume 15, Part 1, page 233.

A 22487/
Distribution:/
Availability Codes

Dist Avail and/or Special

A-1